

**Chitin Enzymology 2001**

R.A.A. Muzzarelli, ed.

Atec, Italy, 2001

ISBN 88-86889-06-2

# **Conservation of structural elements and catalytic mechanism in the chitinolytic enzymes from *Serratia marcescens***

Gali Prag, Constantinos E. Vorgias\*  
and Amos B. Oppenheim

Department of Molecular Genetics and Biotechnology,  
The Hebrew University-Hadassah Medical School,  
Jerusalem 91120 Israel

\*National and Kapodistrian University of Athens, Faculty of Biology,  
Department of Biochemistry-Molecular Biology, Panepistimiopoli,  
Zographou, 15784 Athens, Greece  
ao@md.huji.ac.il.

## **Introduction**

Chitin degrading enzymes are ubiquitous and are found in prokaryotic and eukaryotic organisms as well as in archaea (Perrakis *et al.*, 1993; Tanaka *et al.*, 1999; Brumberg *et al.*, 2001). *Serratia marcescens* has been used as a model system for the utilization of chitin as a carbon source. In the presence of chitin, *S. marcescens* expresses chitinase A, chitinase B, chitinase C and chitobiase that degrade chitin to *N*-acetylglucosamine (NAG) (Warren, 1996). In addition these cells also express a chitin binding protein (CBP21) with unknown function (Folders *et al.*, 2000).

The structures of *S. marcescens* chitinolytic enzymes, chitinase A, chitinase B and chitobiase were solved and the catalytic domains were found to possess an  $\alpha/\beta$  TIM barrel structure (Perrakis *et al.*, 1994; Tews *et al.*, 1996; van Aalten *et al.*, 2000). In contrast, very little is known about the function and structure of chitinase C. Chitinase A, B and C were assigned to glycosyl hydrolase protein family 18, while chitobiase belongs to protein family 20 (Henrissat, 1991). In these enzymes the active site is

located at the carboxy-terminal end of the  $\beta$ -strands of the  $\alpha/\beta$ -barrel suggesting that these proteins, that greatly differ in their primary sequences, all have a common origin.

Information on the co-crystal structure of chitinase-substrate complexes is scant. Earlier attempts to obtain the co-crystal structures of oligoNAG-chitinase complexes in which the non-cleaved substrate occupies the catalytic site proved unsuccessful (Perrakis *et al.*, 1994; Terwisscha-van-Scheltinga *et al.*, 1996). Creation of mutations of the catalytic residues of *S. marcescens* chitinase A and chitobiase allowed us to obtain co-crystals with the native substrate (Prag *et al.*, 2000). Our investigations led us to suggest that *S. marcescens* enzymes participating in chitin metabolism are all employing a similar catalytic mechanism.

## Results and Discussion

### Sequence conservation at the active site

Multiple alignments of glycosyl hydrolases protein families 18 and 20, show the conservation of residues in the  $\beta$ 4-loop #4; DXDXDXE in family 18 and HXGGDE in family 20. Both signatures include Glu315 of chitinase A and Glu540 of chitobiase that act as the proton donor (Henrissat, 1991; Perrakis *et al.*, 1994; Tews *et al.*, 1996), (Figure 1).

### Structural conservation at the active site

We have mutated the conserved Asp and Glu residues mentioned above of both chitinase A and chitobiase and found that both residues are essential for the catalytic activity. Moreover these mutants allowed us, to determine the crystal structures of chitinase A and chitobiase complexed with their native substrate at the active site (Prag *et al.*, 2000; Papanikolaou, Y.; G. P.; Tavlas. G.; C. E. V.; A. B. O. and Petratos K., in preparation). The structure of the catalytic residues Asp and Glu are similar in both enzymes (Figure 2). However, only in chitinase A, Asp313 appears, in the wild-type, in two alternative conformations Papanikolaou, Y.; G. P.; Tavlas. G.; C. E. V.; A. B. O. and Petratos K., in preparation). Comparison of the structures of the enzyme-substrate complexes show high similarity in the position of the -1 to +1 sugars. It was also observed that in both enzyme-substrate complexes, the planes of the sugars at -1 and +1 are tilted around the scissile bond in a similar manner. In both enzymes a critical chair to boat sugar conformational change is involved in the bending and rotation of the substrate upon binding (Figure 2). These energetically non-favored


structures may favor the hydrolysis reaction. Furthermore, the distance from the proton donor and the glycosidic oxygen is conserved. These findings suggest that chitinase A, chitinase B and chitobiase possess similar catalytic sites.

Family 20		Family 18	
CHB_SERMA	529 QPIKTWHFGGDEAK	542	* CHIA_SERMA 307 FDGVDIDWEFP 317
HEX_STRPL	303 TPGRYLHIGGDEAH	316	* EBAG_STRPL 124 LDGVDFDDEYA 134
HEXA_HUMAN	312 FPDFYLHLGGDEVD	325	* EBA1_FLAME 120 LDGVFFDDEYS 130
HEXB_HUMAN	344 FPDQFIHLGGDEVE	357	* CHLY_HEVBR 119 LDGIDFDIEHG 129
HEX_VIBVU	508 QPLTDYHIGADETA	521	* CHIB_SERMA 136 FDGVDIDWEYP 146
CHB_VIBHA	526 APLTTWHFGGDEAK	539	CHIC_SERMA 133 FDGLDIDLEQA 143
HEXB_MOUSE	323 FPDQFIHLGGDEVE	336	CHID_BACCI 295 FNGLDIDLEGS 305
HEXA_MOUSE	312 FPDFYLHLGGDEVD	325	CHIX_STROI 376 LDGLDIDFEGH 386
HEXB_ALTZO	520 VPLNTYHIGADETA	533	CHI1_APHAL 163 FDGIDIDWEYP 173
HEXB_FELCA	286 FPDHFVHLGGDEVE	299	CHI1_BACCI 196 FDGVLDLWEYP 206
HEX1_CANAL	313 FIDDFVHVGNDLQ	326	CHI1_CANAL 139 VDGDFDFDIENK 149
HEX1_ENTHI	290 FGTDYVHVGGEVW	303	CHI1_COCIM 163 FDGIDIDWEYP 173
HEXA_DICDI	297 FIDNYFHTGGDELV	310	CHI1_RHIOL 158 IDGVLDLIEGG 168
HEXC_BOMMO	359 ESTDMFHMGGDEVS	372	CHI2_COCIM 167 VDGDFDFDIEKG 177
HEXA_PORGI	324 FPGTYFHIGGDECP	337	CHI3_CANAL 149 LDGFDFDIENN 159
STRH_STRPN	349 KKTEIFNIGLDEYA	362	CHI4_TRIHA 163 FDGIDIDWEYP 173
STRH_STRPN	793 GKTKIFNFGTDEYA	806	CHIA_ALTZO 305 YDGVLDIDWEFP 315
			CHIB_TOBAC 143 LDGIDFDIELG 153
			CHIT_CAEEL 171 FDGIDIDWEYP 181
			DIAC_HUMAN 135 MDGINIDIEQE 145
			CHIT_YEAST 149 VDGDFDFDIENN 159
			** C3L1_HUMAN 131 FDGLDLAWLYP 142
			** OGP_HUMAN 133 FDGLDLFFLYP 143
			** NARBONIN 124 IDGIDIHYEHI 134
			** CONB_CANEN 123 LDGIHFIDIQKP 133

**Figure 1.** Sequence alignment of protein families 18 and 20 showing the conserved signatures. \* Marks proteins whose structure was solved; \*\* proteins with no chitinase activity. Catalytic glutamic residues are shown in red and the conserved aspartic residues in blue.

The sequences of proteins belonging to glycosyl hydrolase families 18 and 20 are too different to be aligned on the basis of their primary

sequences only. However, the structure of the catalytic  $\alpha/\beta$ -barrel domain is highly conserved. We found that the structure of residues in the conserved motives (DE of chitobiase and DWE of chitinase A) together with the structure of the -1/+1 diNAG is sufficient to obtain structural alignment of chitobiase with chitinase A. Chitinase B could be aligned with chitinase A as these proteins are very similar (Figure 3). This analysis revealed, for example, a conserved tyrosine residue (Tyr669 of chitobiase and Tyr390 of chitinase A). Mutations Tyr669Asp and Tyr390Phe showed reduced activity.




**Figure 2.** Conservation of chitinase A and chitobiase catalytic sites. The catalytic residues of chitinase A (left) and chitobiase (right) are shown and the overlap of the two enzymes (right). The -1 and +1 sugars are shown. Distances are given in Å. The complex of wild-type chitinase A with the substrate was modeled based on mutant complexes.

### Catalytic mechanism

In general, glycosidases degrade carbohydrates by a general acid-base catalysis that involves two amino acid residues, a proton donor and nucleophile. Hydrolysis of the scissile bond results in either the retention or the inversion of the C1-carbon anomeric configuration. While the catalytic glutamate is highly conserved, no amino acid residue that could act as a nucleophile was identified in protein families 18 and 20. A different catalytic mechanism was therefore proposed. Based on the complex of hevamine (a family 18 plant chitinase) with allosamidin and on the complex of chitobiase with diNAG it was suggested that catalysis take


place via substrate-assisted mechanism (Figure 4), (van Scheltinga *et al.*, 1994; Tews *et al.*, 1996 and Prag *et al.*, 2000). In this proposed mechanism the glutamate residue acts as a proton donor, while the terminal oxygen (O7) of the acetamido group of the -1 NAG acts as a nucleophile. In order to act as a nucleophile the acetamido group has to rotate around the C2-N2 bond as was experimentally observed (Prag *et al.*, 2000). However, direct evidence for substrate-assisted catalysis is not yet available.



**Figure 3.** Structural comparison of the TIM-barrel domain of chitinase A (A), chitinase B (B), chitobiase (C) and superposition of all enzymes (D).

What is then the function of the essential aspartic residue in these two protein families? Structural analysis of alanine replacement mutations in chitobiase and chitinase A shed some light on this question. In chitobiase, the complex of Asp539Ala-diNAG the acetamido group is flipped by about  $180^{\circ}$  with respect to that found in the wild-type and in the Glu540Asp complexes (Prag *et al.*, 2000). In this position the acetamido group cannot participate in the substrate-assisted reaction. Thus we suggested that one function of Asp539 is to ensure placing the acetamido group in the conformation that favors the catalysis via the substrate assisted catalysis. In addition, Asp539 appears to be required for restraining the movement of Glu540. Asp539 may provide additional negative charge at the active site

and stabilize the partial positive charge of the acetamido group while forming the oxazoline ring.



**Figure 4.** Model for the catalytic mechanism of chitinase A and chitobiase focusing on the catalytic Asp and Glu residues at the active site.

The analysis of a number of chitinase A-substrate complexes suggests that Asp313 play a similar role in ensuring the positioning of the acetamido group in the conformation that favors the catalysis via the substrate-assisted reaction. Furthermore, we identified in improved crystal structure of wild-type chitinase A that Asp313 is found in two alternative conformations, suggesting that the flexibility of Asp313 is an essential part of the catalytic reaction.


### **Evolutionary consideration**

This study provides further information on the possible evolution of the pathway for chitin metabolism in which the catalytic mechanism and key catalytic residues are conserved. The large differences in amino acid sequence and the presence of additional domains led to the suggestion that proteins possessing an  $\alpha/\beta$  TIM-barrel domain evolved by convergent evolution (Branden & Tooze, 1991; Perrakis *et al.*, 1996). Our analysis lead us to favor the hypothesis that the genes coding for protein families 18 and 20 diverged from a common ancestral gene coding for a TIM-barrel domain. These diverge to acquire different substrate specificity. Evolutionary forces, however, conserved loop #4 with its catalytic glutamate residue. Further tinkering led to the establishment of the signatures around the catalytic sites while conserving the Asp-Glu structure required for catalysis. Interestingly, family 19 chitinases, that do not possess a TIM-barrel structure, utilize an alternative, acid-base catalytic mechanism (Hart *et al.*, 1995). We anticipate that it will be possible to convert the specificity of families 18 and 20 enzymes by modifying their substrate binding properties without altering the catalytic residues (Altamirano *et al.*, 2000).

Several members of family 18, such as the plant seed storage proteins concanavalin B and narbonin, are known to be devoid of catalytic activity (Figure 1, Hennig *et al.*, 1995; Hennig *et al.*, 1992). The structure of the inactive catalytic domain of these proteins is similar to chitinase A and other chitinases. In both inactive proteins, the conserved motif is modified to DIQ in concanavalin B and HYE in narbonin (Terwisscha-van-Scheltinga *et al.*, 1996). Similarly, two human proteins, C3L1 and OGP39, were recently identified as inactive family 18 proteins. Our model for the function of Asp313 and Glu315 of chitinase A provides a molecular explanation for the lack of catalytic activity of these proteins.

### **Cooperative action in chitin degradation**

The biochemical and structural information presented here provide a more complete picture to explain how *S. marcescens* employs its enzymes in chitin degrading. Chitin degradation is probably initiated by the action of endochitinases, probably chitinase A and chitinase C (Watanabe *et al.*, 1997), (Figure 5). Chitinase A also acts as an exochitinase, cleaving diNAG dimers from the reduced end. Chitinase B acts as an exochitinase



**Figure 5.** The cooperative action of chitinases and chitobiase in chitin degradation.

cleaving triNAG and diNAG from the non-reducing end of the chitin oligoNAG chains generated by the action of chitinase A and C (Brurberg *et al.*, 1996; van Aalten *et al.*, 2000). The oligomers, triNAG and diNAG, are subsequently degraded to metabolizable NAG monomers by chitobiase. This analysis clearly explains the synergistic activity of these enzymes that has been reported previously (Brurberg *et al.*, 1996; Watanabe *et al.*, 1999).

### Conclusion

Developments in solving the detailed structures of the chitinolytic enzyme-substrate complexes are expected to result in deeper understanding of the mechanism of action of these enzymes. The accumulated genetic, biochemical and structural data lead us to propose that both chitinase A and chitobiase share a similar catalytic mechanism. This conclusion is based on the following key findings:

- The catalytic domains assume a TIM-barrel structure in which a glutamate residue, located in loop # 4, acts as a proton donor.
- The spatial organization of the conserved Asp and Glu catalytic residues is conserved.



- The structure of the sugar residues where cleavage is to take place is highly conserved.

Furthermore, a model of how *S. marcescens* degrades chitin was developed.

### Acknowledgments

We wish to thank Dr. Vincent Eijsink for valuable critical reading of the manuscript. This work was supported by research grants from the European Commission (EC) BIOTECH 4 Biotechnology and a short term EMBO fellowship (to G.P.). This work was performed, in part, in the Irene and Davide Sala Laboratory for Molecular Genetics.

### References

- Altamirano, M. M., Blackburn, J. M., Aguayo, C. & Fersht, A. R. (2000). Directed evolution of new catalytic activity using the alpha/beta-barrel scaffold. *Nature* 403, 617.
- Branden, C. & Tooze, J. (1991). Introduction to protein structure, Garland Publishing, Inc., New York.
- Brurberg, M. B., Nes, I. F. & Eijsink, V. G. H. (1996). Comparative studies of chitinases A and B from *Serratia marcescens*. *Microbiology* 142, 1581.
- Brurberg, M. B., Synstad, B., Klemsdal, S. S., van Aalten, D. M. F., Sundheim, L. & Eijsink, V. G. H. (2001). Chitinases from *Serratia marcescens*. *Recent Research Developments in Microbiology* (In Press).
- Folders, J., Tommassen, J., van Loon, L. C. & Bitter, W. (2000). Identification of a chitin-binding protein secreted by *Pseudomonas aeruginosa*. *J Bacteriol* 182, 1257.
- Hart, P. J., Pflugger, H. D., Monzingo, A. F., Hollis, T. & Robertus, J. D. (1995). The refined crystal structure of an endochitinase from *Hordeum vulgare* L. seeds at 1.8 Å resolution. *J Mol Biol* 248, 402.
- Hennig, M., Jansonius, J. N., Terwisscha-van-Scheltinga, A. C., Dijkstra, B. W. & Schlesier, B. (1995). Crystal structure of concanavalin B at 1.65 Å resolution. An "inactivated" chitinase from seeds of *Canavalia ensiformis*. *J Mol Biol* 254, 237.
- Hennig, M., Schlesier, B., Dauter, Z., Pfeffer, S., Betzel, C., Hohne, W. E. & Wilson, K. S. (1992). A TIM barrel protein without enzymatic activity? Crystal-structure of narbonin at 1.8 Å resolution. *Febs Lett* 306, 80.
- Henrissat, B. (1991). A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem J* 280, 309.
- Perrakis, A., Tews, I., Dauter, Z., Oppenheim, A. B., Chet, I., Wilson, K. S. & Vorgias, C. E. (1994). Crystal structure of a bacterial chitinase at 2.3 Å resolution. *Structure* 2, 1169.
- Perrakis, A., Tews, I., Wilson, K. S. & Constantin, V. E. (1996). Structural aspects on the catalytic mechanism of chitinases, hevamine, and chitobiasis "far away and yet so close?". In *Chitin Enzymology* (Muzzarelli, R. A. A., ed.), Vol. 2, p. 109. Atec, Italy.

Perrakis, A., Wilson, K. S., Oppenheim, A. B., Chet, I. & Constantin, V. E. (1993). Phylonetic relationships of chitinases. In *Chitin Enzymology* (Muzzarelli, R. A. A., ed.), Vol. 1, p. 232. Atec, Italy.

Prag, G., Papanikolaou, Y., Tavlas, G., Vorgias, C. E., Petratos, K. & Oppenheim, A. B. (2000). Structures of chitobiase mutants complexed with the substrate Di-N-acetyl-d-glucosamine: the catalytic role of the conserved acidic pair, aspartate 539 and glutamate 540. *J Mol Biol* 300, 611.

Tanaka, T., Fujiwara, S., Nishikori, S., Fukui, T., Takagi, M. & Imanaka, T. (1999). A unique chitinase with dual active sites and triple substrate binding sites from the hyperthermophilic archaeon *Pyrococcus kodakaraensis* KOD1. *Appl Environ Microbiol* 65, 5338.

Terwisscha-van-Scheltinga, A. C., Hennig, M. & Dijkstra, B. W. (1996). The 1.8 Å resolution structure of hevamine, a plant chitinase/lysozyme, and analysis of the conserved sequence and structure motifs of glycosyl hydrolase family 18. *J Mol Biol* 262, 243.

Tews, I., Perrakis, A., Oppenheim, A., Dauter, Z., Wilson, K. S. & Vorgias, C. E. (1996). Bacterial chitobiase structure provides insight into catalytic mechanism and the basis of Tay-Sachs disease. *Nat Struct Biol* 3, 638.

van Aalten, D. M. F., Synstad, B., Brurberg, M. B., Hough, E., Riise, B. W., Eijsink, V. G. H. & Wierenga, R. K. (2000). Structure of a two-domain chitotriosidase from *Serratia marcescens* at 1.9-Å resolution. *PNAS* 97, 5842.

Warren, R. A. J. (1996). Microbial hydrolysis of polysaccharides. *Annu. Rev. Microbiol.* 50, 183.

Watanabe, T., Kimura, K., Sumiya, T., Nikaidou, N., Suzuki, K., Suzuki, M., Taiyoji, M., Ferrer, S. & Regue, M. (1997). Genetic analysis of the chitinase system of *Serratia marcescens* 2170. *J Bacteriol* 179, 7111.